THREE-DIMENSIONAL MODELING OF SHOCK WAVE FORMATION AND THROWING OF METAL SHELLS BY HIGH-ENERGY COMPOUNDS

V.A. Shargatov and S.V. Gorkunov

National Research Nuclear University MEPhI, 31 Kashirskoye Shosse, Moscow 115409, Russian Federation

Abstract: The problem of destruction and throwing of a metal shell with the formation of shock waves as a result of the rapid decomposition of a high-energy compound is considered. Mathematical models are formulated to describe the main phenomena, a method for numerical solution is developed taking into account the specific features of flows at different stages of the development of the process. Computational procedures are implemented in the form of program codes intended for performing calculations on high-performance systems. It is shown that the calculation results are in good agreement with the experimental results.

Keywords: numerical simulation; formation of shock waves; throwing of a metal shell

DOI: 10.30826/CE21140210

Figure Captions

Figure 1 Schematic of a high explosive in a shell

Figure 2 Scheme of prototype placement

Figure 3 Prototype geometry. Dimensions are in millimeters

Figure 4 Pressure fields at different moments of time: (a) t = 0.015 ms; (b) 0.03; and (c) t = 0.08 ms

Figure 5 Pressure fields for three instants of time preceding the restructuring of the grid. Computational domain size: (a) $1 \times 1 \times 1$ m; (b) $1.6 \times 1.6 \times 1.6$; and (c) $2.4 \times 2.4 \times 2.4$ m

Figure 6 Pressure profiles at a distance of 1 (a), 2 (b), 2.8 (c), and 5.8 m (d): 1 - experiment; and 2 - simulation

Acknowledgments

The work was supported by the Ministry of Science and Higher Education of the Russian Federation (Government Order Project No. 0723-2020-0036).

References

- Balakrishnan, K., A. L. Kuhl, J. B. Bell, and V. E. Beckner. 2012. An empirical model for the ignition of explosively dispersed aluminum particle clouds. *Shock Waves* 22:591– 603. doi: 10.1007/s00193-012-0388-5.
- 2. Lee, E. L., and C. M. Tarver. 1980. Phenomenological model of shock initiation in heterogeneous explosives. *Phys. Fluids* 23(12):2362.
- Kuhl, A. L., and B. Khasainov 2007. Quadratic model of thermodynamic states in SDF explosions. 38th Conference (International) of ICT Energetic Materials. Karlstruhe, Germany.
- 4. Monaghan, J. J. 1988. An introduction to SPH. *Comput. Phys. Commun.* 48:89–96.
- Parshikov, A. N., and S. A. Medin. 2002. Smoothed particle hydrodynamics using interparticle contact algorithms. *J. Comput. Phys.* 180(1):358–382.

- 6. Kolgan, V. P. 1972. Primenenie printsipa minimal'nykh znacheniy proizvodnoy k postroeniyu konechnoraznostnykh skhem dlya rascheta razryvnykh resheniy gazovoy dinamiki [Application of the principle of minimum derivative values to the construction of finite-difference schemes for calculating discontinuous solutions for gasdynamics]. *Uchenye zapiski TSAGI* [TsAGI Science J.] 3(6):68–77.
- Shargatov, V. A., A. S. Pecherkin, A. S. Sofin, *et al.* 2018. Modeling of shock wave propagation over the obstacles using supercomputers. *J. Phys. Conf. Ser.* 1099(1):012014.
- 8. Ohrt, P.O., and S.J. Alan. 2006. Measured airblast environment from an explosive charge having a scored metal casing. *Symposium (International) on the Military Aspects of Blast and Shock.*
- 9. Victorov, S. B., S. A. Gubin, I. V. Maklashova, and I. I. Revyakin. 2001. Thermodynamic TDS code: Application to

GORENIE I VZRYV (MOSKVA) - COMBUSTION AND EXPLOSION 2021 volume 14 number 2

detonation properties of condensed explosives. *Energetic Materials, Ignition, Combustion and Detonation: 32nd Annu-*

al Conference (International) of ICT Proceedings. Karlsruhe, Germany. 69/1–69/15.

Received May 14, 2021

Contributors

Shargatov Vladimir A. (b. 1958) — Candidate of Science in physics and mathematics, leading research scientist, National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), 31 Kashirskoe Sh., Moscow 115409, Russian Federation; shargatov@mail.ru

Gorkunov Sergey V. (b. 1992) — research engineer, National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), 31 Kashirskoe Sh., Moscow 115409, Russian Federation; gorkunov.ser@mail.ru